




























































































Manufacturer	Species	Brand Name	Prescription	External Parasites					Internal Parasites					Active Ingredient(s)	Application	Duration	Minimum Age (Minimum Weight)		Important Notes
				Fleas	Ticks	Lice	Mites	Biting Flies	Heartworm	Hookworm	Roundworm	Whipworm	Tapeworm				In Cats	In Dogs	
Zoetis	Cats and Dogs	Revolution 	R			-		-				-	-	selamectin	Topical	30 days	8 weeks	6 weeks	Selamectin belongs to a class of medications called macrocyclic lactones. Dogs with MDR1 sensitivity (commonly found in herding breeds like Collies) are at higher risk of toxicity, if using this medication alongside other macrocyclic lactone drugs.
	Cats Only	Revolution Plus 	R			-		-				-	-	sarolaner, selamectin	Topical	30 days	8 weeks (2.8 lbs)	n/a	Sarolaner belongs to a class of medications called isoxazolines. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Medications with this active ingredient should be used with caution in pets with a history of seizures or neurologic disorders. Selamectin belongs to a class of medications called macrocyclic lactones. Dogs with MDR1 sensitivity (commonly found in herding breeds like Collies) are at higher risk of toxicity, if using this medication alongside other macrocyclic lactone drugs.
	Dogs Only	Simparica 	R			-	-	-	-	-	-	-	-	sarolaner	Oral	30 days	n/a	8 weeks (2.8 lbs)	Sarolaner belongs to a class of medications called isoxazolines. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Medications with this active ingredient should be used with caution in pets with a history of seizures or neurologic disorders.
	Dogs Only	Simparica Trio 	R			-	-	-				-	-	sarolaner, moxidectin, pyrantel	Oral	30 days	n/a	8 weeks (2.8 lbs)	Sarolaner belongs to a class of medications called isoxazolines. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Medications with this active ingredient should be used with caution in pets with a history of seizures or neurologic disorders.
Bayer	Cats and Dogs	Advantage II 				-	-	-	-	-	-	-	-	imidacloprid, pyriproxyfen	Topical	30 days	8 weeks (2lbs)	7 weeks (3lbs)	This medication is applied topically, and its active ingredients provides protection against fleas at the level of skin and hair. Bathing may reduce its effectiveness.
	Cats and Dogs	Advantage Multi 				-						-	-	imidacloprid, moxidectin	Topical	30 days	8 weeks (2lbs)	7 weeks (3lbs)	This medication is applied topically, and it's active ingredient imidacloprid provides protection against fleas solely at the level of skin and hair. Bathing may reduce its effectiveness.
	Dogs only	K9 Advatnix II 						-	-	-	-	-	-	imidacloprid, permethrin, pyroproxifen	Topical	30 days	n/a	7 weeks (4lbs)	DO NOTE USE IN CATS. Permethrin is extremely toxic to cats. This medication is applied topically, and its active ingredients provides protection against fleas at the level of skin and hair. Bathing may reduce its effectiveness.
	Dogs only	Multi Duo 	**Dual-pack that includes Advantage Multi and K9 Advantix II**											imidacloprid, moxidectin, permethrin, pyroproxifen	Topical	30 days (Separate Advantage Multi from K9 Advantix II by 14 days)	n/a	7 weeks (4lbs)	DO NOTE USE IN CATS. Permethrin is extremely toxic to cats. These medications are applied topically, and their active ingredients imidacloprid, permethrin and pyroproxifen provide protection against parasites at the level of skin and hair. Bathing may reduce its effectiveness.
Novartis	Cats and Dogs	Capstar 			-	-	-	-	-	-	-	-	-	nitentpyram	Oral	Every 7 days, until adult fleas are no longer observed	4 weeks (2lbs)	4 weeks (2lbs)	
Merck	Cats and Dogs	Bravecto 	R			-	-	-	-	-	-	-	-	fluralaner	Topical (cats and dogs), Oral (dogs only)	90 days	6 months (2.6lbs)	6 months (4.4lbs)	Fluralaner belongs to a class of medications called isoxazolines. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Medications with this active ingredient should be used with caution in pets with a history of seizures or neurologic disorders.
Elanco	Cats and Dogs	Interceptor 	R	-	-	-	-					-	-	milbemycin	Oral	30 days	6 weeks (1.5 lbs)	4 weeks (2lbs)	Milbemycin belongs to a class of medications called macrocyclic lactones. Dogs with MDR1 sensitivity (commonly found in herding breeds like Collies) are at higher risk of toxicity, if using this medication alongside other macrocyclic lactone drugs.
	Dogs only	Interceptor Plus 	R	-	-	-	-						-	milbemycin, praziquantel	Oral	30 days	n/a	6 weeks (2lbs)	Milbemycin belongs to a class of medications called macrocyclic lactones. Dogs with MDR1 sensitivity (commonly found in herding breeds like Collies) are at higher risk of toxicity, if using this medication alongside other macrocyclic lactone drugs.
Boehringer Ingelheim	Cats Only	Nexgard Combo 	R			-		-						esafoxolaner, eprinomectin, praziquantel	Topical	30 days	8 weeks (1.8 lbs)	n/a	Esafoxolaner belongs to a class of medications called isoxazolines. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Medications with this active ingredient should be used with caution in pets with a history of seizures or neurologic disorders. Milbemycin belongs to a class of medications called macrocyclic lactones. Dogs with MDR1 sensitivity (commonly found in herding breeds like Collies) are at higher risk of toxicity, if using this medication alongside other macrocyclic lactone drugs.
	Dogs Only	Heartgard Plus 	R	-	-	-	-				-	-	ivermectin, pyrantel	Oral	30 days	n/a	6 weeks	Ivermectin belongs to a class of medications called macrocyclic lactones. Dogs with MDR1 sensitivity (commonly found in herding breeds like Collies) are at higher risk of toxicity, if using this medication alongside other macrocyclic lactone drugs.	
	Dogs Only	Nexgard 	R			-	-	-	-	-	-	-	-	afoxolaner	Oral	30 days	n/a	8 weeks (4lbs)	Afoxolaner belongs to a class of medications called isoxazolines. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Medications with this active ingredient should be used with caution in pets with a history of seizures or neurologic disorders.
	Dogs Only	Nexgard Spectra 	R			-	-	-					-	afoxolaner, milbemycin	Oral	30 days	n/a	8 weeks (4.4 lbs)	Afoxolaner belongs to a class of medications called isoxazolines. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Medications with this active ingredient should be used with caution in pets with a history of seizures or neurologic disorders. Milbemycin belongs to a class of medications called macrocyclic lactones. Dogs with MDR1 sensitivity (commonly found in herding breeds like Collies) are at higher risk of toxicity, if using this medication alongside other macrocyclic lactone drugs.

Legend

-  Treats indicated parasite spp. in cats
-  Treats indicated parasite spp. in dogs
-  Treats indicated parasite spp. in cats and dogs
- R** A prescription is required to get this medication